

Assessing delivery of the five 'As' for patient-centered counseling

RUSSELL E. GLASGOW, SETH EMONT¹ and DORIANE C. MILLER²

Clinical Research Unit, Kaiser Permanente Colorado, Denver, CO, USA, ¹White Mountain Research Associates, L.L.C., Danbury, NH, USA and ²Rush Medical College, Chicago, IL, USA

SUMMARY

The '5As' model of behavior change provides a sequence of evidence-based clinician and office practice behaviors (Assess, Advise, Agree, Assist, Arrange) that can be applied in primary care settings to address a broad range of behaviors and health conditions. Although the 5As approach is becoming more widely adopted as a strategy for health behavior change counseling, practical and standardized assessments of 5As delivery are not widely available. This article provides clinicians and researchers with alternatives for assessment of 5As implementation for both quality improvement, and for research and evaluation

purposes, and presents several practical tools they may wish to use. Sample instruments for tracking delivery of the 5As and related tools that are in the public domain are provided to facilitate integration of self-management support into clinical care. We discuss the strengths and limitations of the various assessment approaches. Promising and practical measures to assess the 5As exist for both quality improvement and research purposes. Additional validation is needed on almost all current procedures, and both clinicians and researchers are encouraged to use these instruments and share the resulting data.

Key words: assessment; primary care; behavior change

INTRODUCTION

The '5As' model of behavior change counseling is an evidence-based approach appropriate for a broad range of different behaviors and health conditions, and is feasible to apply in primary care (Fiore *et al.*, 2000; Glasgow *et al.*, 2001a; Glasgow *et al.*, 2002; Serdula *et al.*, 2003; Glasgow *et al.*, 2004b; Goldstein *et al.*, 2004; The Quality Indicator Study Group, 1995). The 5As are as follows: *assessing* patient level of behavior, beliefs and motivation; *advising* the patient based upon personal health risks; *agreeing* with the patient on a realistic set of goals; *assisting* to anticipate barriers and develop a specific action plan; and *arranging follow-up* support (Moen *et al.*, 1999; Glasgow *et al.*, 2002; Berwick, 2003; Glasgow *et al.*, 2003).

However, there are no standard, widely used assessments of 5As delivery. Most studies of

the 5As have developed their own measures. As the 5As become more widely adopted, it is increasingly important to have reliable, valid and practical measures of these strategies.

Recently, the Lifescripts consortium, through funding by the Australian Government Department of Health and Ageing, released a series of practice tools and other resources to support general practitioners in reducing lifestyle risk factors among their patients (National Heart Foundation of Australia and Kinect Australia for the Lifescripts Consortium, 2005). The Lifescripts package contains a variety of evidence-based resources, manuals and user-friendly assessment tools for GPs, practice nurses and staff in the general practice setting. The program and associated assessment tools build on the SNAP framework developed by the Royal

Australian College of General Practitioners (Smoking, Nutrition, Alcohol and Physical activity) (The Royal Australian College of General Practitioners, 2004). This framework utilizes the 5As for incorporating effective clinical strategies for the detection, assessment and management of SNAP risk factors in general practice settings.

Documentation and tracking of the 5As as a way of delivering self-management support is now required by the Joint Commission for the Accreditation of Health Care Organizations as part of its Disease Management Certification program. Health care organizations satisfying this requirement as part of their disease management protocols are more likely to receive certification. This certification could provide health care organizations with leverage to negotiate higher reimbursement rates from third-party payers.

The purposes of this article are to (i) discuss assessment of the 5As from both quality improvement (QI) and research perspectives, (ii) describe and provide examples of instruments that have been used to assess 5As implementation and (iii) discuss the strengths and limitations of present 5As assessment approaches.

There are four primary sources of information that can be used to assess delivery of the 5As as follows: direct observation of patient–clinician encounters; medical records; clinician reports; and patient reports (Table 1). Each has its strengths and limitations, and is used for different purposes. Multiple approaches were used to identify potential instruments as follows: (i) an electronic search was conducted using PubMed

with various combinations of the keywords ‘5As, behavioral counseling, assessment and physician advice’; (ii) recent articles and reviews of behavioral counseling that included the 5As were consulted and reference lists in these articles were searched; and (iii) contacts were made with leading researchers in the area. Given that there are no MeSH headings related to the 5As and the recency of much of the work in this area, the latter two approaches identified the majority of tools. We describe each of the approaches organized by the primary purpose of the assessment as follows: (i) QI or (ii) research and evaluation, since the issues and selection criteria are different.

Quality improvement

To be of value for QI, measures need to be brief, easy and inexpensive to administer and score, and capable of providing useful and ‘real-time’ feedback. QI measures also should allow for comparisons over time and can be benchmarked against data from other clinicians or settings, across populations, or standards of care. Although measurement tracking for QI is complementary in many respects to ongoing data collection for research, a key difference is that measurement tracking for QI has as its primary goal *continuous* practice improvement (i.e. rapid analysis of data, reporting and feedback).

Below, we describe various approaches to tracking delivery of the 5As for QI. These approaches should be practical for integration into routine care in most settings. The various

Table 1: 5As assessment approaches by primary purposes and cost

Approach (references)	Quality improvement		Research		Cost
	Rapid	Longer	Pop-based	Detailed	
Staff completed instruments					
Checklists	XX	X	XX		Low
Surveys		X		X	High
Observational or administrative records					
Recorded interactions		X		XX	Highest
EMR/chart reviews		X	X	XX	High initial Low ongoing
Patient report measures					
Brief checklists	X	X	X		Moderate
Detailed surveys		XX		XX	High
Satisfaction with care	X	X	X		Low ongoing
Prompts and print-outs	X	X	X		High initial Low ongoing

X = usually associated with this approach; XX = key characteristics of this approach.

assessment approaches also can be used in *combination* to provide a more comprehensive picture.

Checklists are designed to both prompt and record clinician behaviors. Out of the four approaches described, this is perhaps the most efficient, since it is a 'real-time' approach to

documenting patient-clinician interactions. Checklists can be included in the patient record and provide the clinician with a framework to record follow-up notes. Figure 1 is a checklist that was recently developed for the Health Research & Educational Trust's Pilot Collaborative on Self-Management Support

PATIENT NAME _____ DATE _____ TYPE CONTACT: _____

(Check ALL that apply)

___ ASSESS- Patient completed assessment and received feedback on health behaviors DATE: _____

Notes: _____

___ ADVISE- Personally relevant, specific recommendation for behavior change DATE: _____

Notes: _____

___ AGREE- Document collaboratively set, specific achievable behavioral goal with patient DATE: _____

Notes: _____

___ ASSIST- a. ___ Worked with patient to develop action plan, strategies, or problem solve DATE: _____
 b. ___ Referred patient to resource, counselor, or group to work on above

Notes: _____

___ ARRANGE- a. ___ Made follow-up contact within 2 weeks after setting behavioral goal DATE: _____
 b. ___ Checked to see that patient completed referral visit
 c. ___ Obtained feedback on patient progress from referral party or directly from patient

Notes: _____

GENERAL NOTES: _____

Fig. 1: Delivery of 5As self-management support.

Kaiser Foundation Health Plan of the Northwest Tobacco Checklist			Date	Time	Location	Ordering Clinician
<input type="checkbox"/> Current Smoker <input type="checkbox"/> Recently Quit			Patient Name			
Clinician: <u>Yes</u> <u>No</u> Advice Given <input type="checkbox"/> <input type="checkbox"/> Referred to Nurse <input type="checkbox"/> <input type="checkbox"/> Accepted Referral <input type="checkbox"/> <input type="checkbox"/> Clinician Signature: _____			Chart No.			
Nurse/Clinic Staff: <u>Yes</u> <u>No</u> Patient seen by nurse/clinic staff <input type="checkbox"/> <input type="checkbox"/> Video shown, "When the time is right" <input type="checkbox"/> <input type="checkbox"/> Video shown, "FFC" <input type="checkbox"/> <input type="checkbox"/> Quit date set <input type="checkbox"/> <input type="checkbox"/> Quit date: ___/___/___ Best time to call: <input type="checkbox"/> a.m. <input type="checkbox"/> p.m. <input type="checkbox"/> eve <input type="checkbox"/> weekend Best place: <input type="checkbox"/> home <input type="checkbox"/> work <input type="checkbox"/> no call Signature: _____			Phone Number (include area code)			
			H: W:			
			Comments:			

Fig. 2: Tobacco care checklist. Adapted from: Hollis et al. (Hollis, Bills, Whitlock et al., 2000)

(see www.collaborativeselfmanagement.org). It allows the clinician to ‘walk’ patients through the 5As at a given visit, and to later re-assess progress in meeting collaboratively set goals. A slightly more complex example is shown in Figure 2. This checklist, which embeds 5As issues within counselor responsibilities, was developed specifically for a tobacco use cessation intervention (Hollis *et al.*, 2000). Although the checklist is perhaps the easiest approach to document 5As counseling for QI purposes, clinicians must use it consistently across visits if it is to provide reliable data.

Patient exit interviews assess patient perspectives on the counseling they received, and evaluate changes in practice patterns related to the 5As. Exit interviews can be conducted on a random sample from the patient population or with a consecutive set of patients at prescribed intervals (e.g. monthly or quarterly) to track changes in practice patterns over time. Figure 3 provides 10 questions that can be used to assess patient perceptions of delivery of the 5As. This question set was used by sites participating in the Pilot Collaborative on Self-Management Support to assess delivery of four of the 5As—Advise, Agree, Assist and Arrange. The Self-Management Pilot Collaborative planning team

has also developed a 5-item patient satisfaction survey, which specifically addresses whether the patient thought she received counseling around the 5As, and if so, how satisfied she was with the care received. It can be conducted via patient exit interview or self-administered (Figure 4). Such surveys can be conducted in combination with provider checklists to triangulate clinician-reported delivery of care with patient-reports of care received. This is important, since it is possible for patients to report that they did not receive specific counseling even though the clinician may have reported doing so. From a QI perspective, this could help identify gaps and improvements needed to effectively communicate with patients.

Medical records can be abstracted to document 5As delivery. Although this is a labor-intensive process, medical chart abstraction provides perhaps one of the most important measures of clinicians’ delivery of the 5As. This is because what gets into the medical record is often what other team members see and pay attention to, and such documentation provides a prompt for follow-up on other ‘As’ at future contacts. A limitation of chart audits is that they typically under-report the level of counseling (Stange *et al.*, 1998; Newell *et al.*, 1999). Chart abstraction can be

Tell us what you think about your health care

These questions are about your visits and conversations with your health care team.

Members of your *health care team* are doctors, nurses or any medical person who gives you health care for your long term health problems.

Check one box for each question.

- | | |
|--|--|
| <p>1. My health care team asked me in a conversation how I take care of my health.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> | <p>7. My health care team helped me make a plan that I can use every day to help take care of my health problems.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> |
| <p>2. My health care team asked me in a survey how I take care of my health.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> | <p>8. My health care team helped me plan ahead so that I can take care of my health problems even during hard times.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> |
| <p>3. My health care team gave me a list of things that I can do to improve my health.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> | <p>9. My health care team told me about other people who can help me with my health problems (such as groups, classes, counselors, dieticians and health educators).</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> |
| <p>4. My health care team gave me personalized advice about how I can improve my health.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> | <p>10. My health care team made plans to contact me after a visit to see how I was doing.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> |
| <p>5. My health care team helped me to set specific goals to manage my health problems.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> | |
| <p>6. My health care team asked for my ideas about how I can take care of my health problems.</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know</p> | |

Thank you. We will keep your answers private and confidential!

Fig. 3: Brief 5As patient survey. *Source:* www.collaborativeselfmanagement.org.

conducted on a consecutive or random sample of patients and could occur concurrently with other QI efforts. It does require development of coding manuals and definitions to ensure that charts are reviewed in a consistent manner (Boyle and Solberg, 2004). For example, Serdula *et al.* developed a weight loss counseling tool using the 5As, based on an evidence-based algorithm for the treatment of obesity (Serdula *et al.*,

2003). Criteria used to define delivery of each of the 5As are also available in various publications (Fiore *et al.*, 2000; Glasgow *et al.*, 2003; Glasgow *et al.*, 2004b; Goldstein *et al.*, 2004). DePue *et al.* utilized a chart audit approach in community health centers to evaluate performance rates of the '4As' (Ask, Advise, Assist, Arrange) for tobacco use cessation counseling (DePue *et al.*, 2002). They audited consecutive

How satisfied are you with your health care?

Tell us what you think about your health care and your health care team. Members of your *health care team* are doctors, nurses or any medical person who gives you health care for your long term health problems.

1. My health care team asked about my **ideas and beliefs** when we talked about my health problems and treatment.
 Yes No Don't know
If you chose yes, how satisfied were you with that conversation?
 Not Satisfied
 Satisfied
 Very Satisfied
 Don't know
2. My health care team told me in a clear way how I could help take care of my health problems.
 Yes No Don't know
If you chose yes, how satisfied were you with that conversation?
 Not Satisfied
 Satisfied
 Very Satisfied
 Don't know
3. My health care team talked with me about setting goals to take care of my health problems.
 Yes No Don't know
If you chose yes, how satisfied were you with that conversation?
 Not Satisfied
 Satisfied
 Very Satisfied
 Don't know
4. My health care team helped me to solve problems so that I could meet my goals and improve my health.
 Yes No Don't know
If you chose yes, how satisfied were you with that conversation?
 Not Satisfied
 Satisfied
 Very Satisfied
 Don't know
5. My health care team talked with me about how to get help from my friends, family and community.
 Yes No Don't know
If you chose yes, how satisfied were you with that conversation?
 Not Satisfied
 Satisfied
 Very Satisfied
 Don't know

Thank you. We will keep your answers private and confidential!

Fig. 4: 5As patient satisfaction survey. *Source:* www.collaborativeselfmanagement.org.

records of each health center site, and related quality of care delivery to site level characteristics (e.g. size, location, percentage of providers attending training).

Brief retrospective surveys can help prompt a discussion with patients about goals and creating self-management supports. Self-report measures

obtain the patient perspective on receipt of counseling and overcome some of the limitations of provider reports, but are themselves subject to reporting biases and social desirability (Stange *et al.*, 1998). Various research groups have developed brief, face valid questions for patients about the extent to which they have received 5As

counseling for specific target behaviors (Sciamanna *et al.*, 2002). Glasgow *et al.* reported that primary care patients responding to a national survey reported receiving 5As for smoking cessation more often than for other health-related behaviors such as physical activity or healthy eating (Glasgow *et al.*, 2001a). They also reported that the last two A's—assistance and arranging follow-up support—were reported the least often across target behaviors.

With more sophisticated electronic patient information systems, it is possible to integrate clinician-based prompts and feedback with other interactive technologies for self-management support (Glasgow and Bull, 2001; Bodenheimer and Grumbach, 2003; Glasgow *et al.*, 2004a). Patient surveys are most feasible when electronic registries are available that can provide reminders to enhance self-management supports at the point of contact (Glasgow *et al.*, 2001b; McKay *et al.*, 2001) although even manual reminders can be effective in this regard (Marquez, 2001). Glasgow *et al.* provide an example of a flow diagram that includes prompts for all personnel, including patients, physicians and care managers involved in a typical patient visit (Glasgow *et al.*, 2003). They provide (i) a scenario of what a patient visit might look like that fully integrated self-management support into primary care and (ii) strategies that can be used by various staff members involved with the patient's care before, during and after a visit.

Key characteristics of the QI assessment approaches described above are that they are practical and relatively inexpensive to implement—especially if chart reviews are being done for other purposes (Table 1). Integrating one or more of these into existing QI strategies also can be cost-effective—for example, through a third-party organization like Quality Improvement Organizations (QIOs). Under the direction of Centers for Medicare and Medicaid Services, the QIO program consists of a national network of 53 QIOs responsible for each US state, territory and the District of Columbia. QIOs work with consumers, clinicians, hospitals and other providers to improve care delivery systems, particularly among underserved populations (www.medqic.org).

Finally, the Health Disparities Collaboratives™ offer a number of training manuals and tools available for free download (www.healthdisparities.net). This website offers a wide array of registries, questionnaires, chart audit

tools, flow charts and reports for components of the Chronic Care Model (i.e. clinical information systems, community, decision support, delivery system design, organization of health care and self-management support) for asthma, depression and diabetes. Although some of these QI aids are not specifically organized around the 5As, most do involve assessment, assistance and arranging follow-up.

Research and evaluation

Instruments used for research purposes are typically held to higher standards of reliability and validity than QI measures. They also tend to be longer, more expensive and comprehensive, and usually cannot be scored quickly to provide immediate feedback. They are intended to provide a more in-depth understanding of the complexities of care delivery (Table 1).

Direct observation methods

Despite their expense and impracticality in many situations, observational coding systems, based upon live, videotaped or audiotaped clinic–patient interactions are considered the gold standard assessment methods for patient–clinician interactions. To our knowledge, there are no observational coding systems specifically designed for the 5As. There are, however, numerous related systems for coding patient–provider interactions (Roter, 1991). The earliest methods were based on the client-centered counseling approach of Rogers and colleagues and related systems have evolved in the psychotherapy literature to measure constructs such as empathy and percent of talk time for patients versus clinicians (Rogers, 1951; Truax, 1966). More recently, researchers interested in patient-centered interventions have coded patient–staff interactions for key components of motivational interviewing and patient-centered communication style (Amrhein *et al.*, 2003; Baer *et al.*, 2004). Closer to primary care, Stange and colleagues have rated patient–provider interactions for the delivery of preventive services (Stange *et al.*, 2000; Flocke and Stange, 2004; Gilchrist *et al.*, 2004). Results from observational coding systems have contributed greatly to understanding the structure and patterns of patient–clinician communication. These systems can also be of great value for clinician training. Because of their cost, complexity and the lack of real-time

feedback, observational coding systems for the 5As are not widely used.

Computer-based assessments of medical records

Electronic chart abstraction tools and coding manuals are available via the Internet. For example, the RAND organization is currently conducting a 4-year study of the effectiveness of chronic illness care collaboratives that utilize the Chronic Care Model (Wagner, 1998; Cretin *et al.*, 2004) across various patient populations, including those with asthma, depression, congestive heart failure and diabetes. They are tracking key processes of care and have developed quality indicators for each disease. Measures related to the 5As associated with the Chronic Care Model are being tracked (i.e. documentation of counseling, goal setting and development of an action plan) (Cretin *et al.*, 2004). These investigators developed a series of Microsoft® Access-based tools to enter the abstracted data (<http://www.rand.org/health/ICICE>).

Smoking cessation is the area in which the 5As have been used most extensively (Fiore *et al.*, 2000; Hollis *et al.*, 2000; DePue *et al.*, 2002). Kaiser Permanente researchers (Hazlehurst *et al.*, 2005) developed a natural language programming tool to analyze free field text notes in electronic medical records to analyze the extent to which each of the 5As is used for smoking cessation counseling in primary care. They found that while patients were often assessed and advised to quit smoking, seldom were they provided assistance or follow-up support.

Patient surveys

Patient-centered activities such as listening to patients, considering their priorities, developing collaborative goals and eliciting coping suggestions that are congruent with patient values, preferences and social environment are at the heart of the 5As approach. Thus, it is critically important to evaluate patient perceptions of the extent to which they have received such assistance. Although a variety of related instruments have been developed, especially to assess core aspects of primary care and delivery of components of the Chronic Care Model, few survey instruments assess the 5As. Some scales of the ACES (Murray and Safran, 2000) and the Shi *et al.* assessments of patient perspectives of primary care are related to the first three of the 5As (Shi *et al.*, 2001).

Glasgow, Wagner and colleagues reported on the Patient Assessment of Chronic Illness Care (PACIC) (Glasgow *et al.*, 2005a; Glasgow *et al.*, 2005b). This instrument assesses implementation of five activities (patient activation, practice design/decision support, goal setting, problem solving and follow-up) related to the Chronic Care Model that are congruent with the 5As (Glasgow *et al.*, 2005b). Glasgow *et al.* report that assistance in problem solving and arranging follow-up support (the 4th and 5th As) are reported least often by patients (Glasgow *et al.*, 2005a).

The PACIC has recently been modified to include additional items so that it can be scored specifically for each of the 5As (Glasgow *et al.*, 2005b) (www.collaborativeselfmanagement.org). To our knowledge, this 26-item survey (Table 2) is the only instrument of this level of comprehensiveness scored specifically for the 5As. Although this scale worked well in a pilot collaborative and is an extension of the validated 20-item PACIC, to date the 26-item scale and the 5As scoring has only been formally validated in one study with diabetes patients (Glasgow *et al.*, 2005b). Both the 20- and 26-item versions of the PACIC are in the public domain.

CONCLUSIONS

This article reviews instruments that we were able to locate through literature searches and contacts with other colleagues, or developed as part of QI efforts to improve self-management support. Since there are no MeSH terms directly related to the 5As, we cannot be certain that all relevant tools were identified. Also, many of the tools are as yet unpublished. To address this situation, we have provided the tables and figures in this manuscript and posted the tools that are in the public domain on the www.collaborativeselfmanagement.org website.

It is not possible to conclude that any one 5As assessment procedure should be recommended over others. Which instrument is best depends on one's purpose and situational constraints. Different instruments would be selected to definitively identify processes associated with improved outcomes in a multisite clinical trial than for a QI project requiring immediate feedback.

Available literature (Bodenheimer *et al.*, 2002; Bodenheimer and Grumbach, 2003; Glasgow

Table 2: Interactions with your health care team

Staying healthy can be difficult when you have a chronic illness. We would like to learn about the type of help with your condition you get from your health care team. This might include your regular doctor, his or her nurse, or physician's assistant who treats your illness. Your answers will be kept confidential and will not be shared with anyone else

Think about the health care you have received over the past 6 months. (If it is been more than 6 months since you have seen your doctor or nurse, think about your most recent visit.)

Over the past 6 months, when receiving medical care for my chronic illness, I was:

	Almost never	Generally not	Sometimes	Most of the time	Almost always
(1) Asked for my ideas when we made a treatment plan	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(2) Given choices about treatment to think about	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(3) Asked to talk about any problems with my medicines or their effects	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(4) Given a written list of things I should do to improve my health	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(5) Satisfied that my care was well organized	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(6) Shown how what I did to take care of my illness influenced my condition	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(7) Asked to talk about my goals in caring for my illness	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(8) Helped to set specific goals to improve my eating or exercise	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(9) Given a copy of my treatment plan	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(10) Encouraged to go to a specific group or class to help me cope with my chronic illness	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(11) Asked questions, either directly or on a survey, about my health habits	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(12) Sure that my doctor or nurse thought about my values and my traditions when they recommended treatments to me	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(13) Helped to make a treatment plan that I could do in my daily life	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(14) Helped to plan ahead so I could take care of my illness even in hard times	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(15) Asked how my chronic illness affects my life	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(16) Contacted after a visit to see how things were going	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(17) Encouraged to attend programs in the community that could help me	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(18) Referred to a dietitian, health educator or counselor	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(19) Told how my visits with other types of doctors, like the eye doctor or surgeon, helped my treatment	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(20) Asked how my visits with other doctors were going	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(21) Asked what I would like to discuss about my illness at that visit	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(22) Asked how <i>my work, family</i> or social situation related to taking care of my illness	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(23) Helped to make plans for how to get support from my friends, family or community	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(24) Told how important what I do to take care of my illness (e.g. exercise, watch my diet) is for my health	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(25) Helped to set a goal with my doctor or health team member	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(26) Given a form or book in which to help me record the progress I am making on my goals	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅
(27) Placed into a telephone-based support program to help manage my condition	<input type="checkbox"/> ₁	<input type="checkbox"/> ₂	<input type="checkbox"/> ₃	<input type="checkbox"/> ₄	<input type="checkbox"/> ₅

5As scoring template

Ask/Assess—average items: 3, 11, 15, 20, 21, 22.

Advise—average items: 4, 6, 19, 24.

Agree—average items: 1, 2, 7, 8, 25.

Assist—average items: 9, 10, 12, 13, 14, 26.

Arrange—average items: 16, 17, 18, 23, 27.

Total score—average score across all (valid) items answered.

et al., 2003; Hill-Briggs, 2003) suggests that the last two As—assistance in problem solving and arranging follow-up support—are especially important to produce meaningful and lasting behavior change. Unfortunately, they are also the two As that seem to be delivered least often (Glasgow et al., 2001a; Glasgow et al., 2005a). Measures of the 5As should devote special attention to assessment of these components.

We hope that this article will stimulate others to use, validate and refine existing 5As assessment procedures. To our knowledge, at present only the PACIC and the natural language programming methods have supportive validation data. Most of the other tools have face validity and may be more practical, but data are needed on their reliability, validity and especially sensitivity to change. Translation into Spanish and other languages, and validation studies with different ethnic and cultural groups should be a high priority.

ACKNOWLEDGEMENT

Preparation of this paper was supported by a grant from the Robert Wood Johnson Foundation, ‘Pilot Collaborative for Self-Management Support’ to Doriane C. Miller.

Address for correspondence:

Russell E. Glasgow, PhD
Clinical Research Unit
Kaiser Permanente Colorado
335 Road Runner Lane
Penrose
CO 81240
USA
E-mail: russg@ris.net

REFERENCES

- Amrhein, P. C., Miller, W. R., Yahne, C. E., Palmer, M. and Fulcher, L. (2003) Client commitment language during motivational interviewing predicts drug use outcomes. *Journal of Consulting and Clinical Psychology*, **71**, 862–878.
- Baer, J. S., Rosengren, D. B., Dunn, C. W., Wells, E. A., Ogle, R. L. and Hartzler, B. (2004) An evaluation of workshop training in motivational interviewing for addiction and mental health clinicians. *Drug and Alcohol Dependence*, **73**, 99–106.
- Berwick, D. M. (2003) Disseminating innovations in health care. *Journal of the American Medical Association*, **289**, 1969–1975.
- Bodenheimer, T. S. and Grumbach, K. (2003) Electronic technology: a spark to revolutionize primary care? *Journal of the American Medical Association*, **290**, 259–264.
- Bodenheimer, T. S., Lorig, K., Holman, H. and Grumbach, K. (2002) Patient self-management of chronic disease in primary care. *Journal of the American Medical Association*, **288**, 2469–2475.
- Boyle, R. and Solberg, L. I. (2004) Is making smoking status a vital sign sufficient to increase cessation support actions in clinical practice? *Annals of Family Medicine*, **2**, 22–25.
- Cretin, S., Shortell, S. M. and Keeler, E. B. (2004) An evaluation of collaborative interventions to improve chronic illness care: framework and study design. *Evaluation Review*, **28**, 28–51.
- DePue, J. D., Goldstein, M. G., Schilling, A., Reiss, P., Papandonatos, G., Sciamanna, C., et al. (2002) Dissemination of the AHCPR clinical practice guideline in community health centers. *Tobacco Control*, **11**, 239–335.
- Fiore, M. C., Bailey, W. C., Cohen, S. J., Dorfman, S. F., Goldstein, M. G., Gritz, E. R., et al. (2000) *Treating Tobacco Use and Dependence: Clinical Practice Guideline*. U.S. Department of Health and Human Services, Public Health Service, Rockville, MD.
- Flocke, S. A. and Stange, K. C. (2004) Direct observation and patient recall of health behavior advice. *Preventive Medicine*, **38**, 343–349.
- Gilchrist, V. J., Stange, K. C., Flocke, S. A., McCord, G. and Bourguet, C. C. (2004) A comparison of the National Ambulatory Medical Care Survey (NAMCS) measurement approach with direct observation of outpatient visits. *Medical Care*, **42**, 276–280.
- Glasgow, R. E. and Bull, S. S. (2001) Making a difference with interactive technology: considerations in using and evaluating computerized aids for diabetes self-management education. *Diabetes Spectrum*, **14**, 99–106.
- Glasgow, R. E., Eakin, E. G., Fisher, E. B., Bacak, S. J. and Brownson, R. C. (2001a) Physician advice and support for physical activity: results from a national survey. *American Journal of Preventive Medicine*, **21**, 189–196.
- Glasgow, R. E., Hiss, R. G., Anderson, R. M., Friedman, N. M., Hayward, R. A., Marrero, D. G., et al. (2001b) Report of Health Care Delivery Work Group: behavioral research related to the establishment of a chronic disease model for diabetes care. *Diabetes Care*, **24**, 124–130.
- Glasgow, R. E., Funnell, M. M., Bonomi, A. E., Davis, C., Beckham, V. and Wagner, E. H. (2002) Self-management aspects of the improving chronic illness care breakthrough series: implementation with diabetes and heart failure teams. *Annals of Behavioral Medicine*, **24**, 80–87.
- Glasgow, R. E., Davis, C. L., Funnell, M. M. and Beck, A. (2003) Implementing practical interventions to support chronic illness self-management in health care settings: lessons learned and recommendations. *Joint Commission Journal on Quality and Safety*, **29**, 563–574.
- Glasgow, R. E., Bull, S. S., Piette, J. D. and Steiner, J. (2004a) Interactive behavior change technology: a partial solution to the competing demands of primary care. *American Journal of Preventive Medicine*, **27**, 80–87.
- Glasgow, R. E., Goldstein, M. G., Ockene, J. and Pronk, N. P. (2004b) Translating what we have learned into practice: principles and hypotheses for addressing multiple behaviors in primary care. *American Journal of Preventive Medicine*, **27**, 88–101.
- Glasgow, R. E., Wagner, E., Schaefer, J., Mahoney, L., Reid, R. and Greene, S. (2005a) Development and validation of the Patient Assessment of Chronic Illness Care (PACIC). *Medical Care*, **43**, 436–444.

- Glasgow, R. E., Whitesides, H., Nelson, C. C. and King, D. K. (2005b) Use of the Patient Assessment of Chronic Illness Care (PACIC) with diabetes patients: relationship to patient characteristics, receipt of care, and self-management. *Diabetes Care*, **28**, 2655–2661.
- Goldstein, M. G., Whitlock, E. P. and DePue, J. (2004) Multiple health risk behavior interventions in primary care: summary of research evidence. *American Journal of Preventive Medicine*, **27** (2 Suppl), 61–79.
- Hazlehurst, B., Sittig, D. F., Stevens, V. J., Smith, K. S., Hollis, J. F., Vogt, T. M., *et al.* (2005) Using natural language processing to assess tobacco care quality in the EMR. *American Journal of Preventive Medicine*, **29**, 434–439.
- Hill-Briggs, F. (2003) Problem solving in diabetes self-management: a model of chronic illness self-management behavior. *Annals of Behavioral Medicine*, **25**, 182–193.
- Hollis, J. F., Bills, R., Whitlock, E. and Stevens, V. J. (2000) Implementing tobacco interventions in the real world of managed care. *Tobacco Control*, **9** (Suppl 1), 18–24.
- Marquez, L. (2001) *Helping Healthcare Providers Perform According to Standards*. Published for the U.S. Agency for International Development (USAID) by the Quality Assurance Project, Bethesda, MD, Operations Research Issue Paper 2(3).
- McKay, H. G., King, D., Eakin, E. G., Seeley, J. R. and Glasgow, R. E. (2001) The Diabetes Network Internet-based physical activity intervention: a randomized pilot study. *Diabetes Care*, **24**, 1328–1334.
- Moen, R. D., Nolan, T. W. and Provost, L. P. (1999) *Quality Improvement Through Planned Experimentation*, 2nd edition. McGraw Hill, New York City.
- Murray, A. and Safran, D. G. (2000) The Primary Care Assessment Survey: a tool for measuring, monitoring, and improving primary care. In Maruish, M. E. (ed.), *Handbook of Psychological Assessment in Primary Care Settings*. Lawrence Erlbaum Associates, Mahwah, NJ, pp. 623–651.
- National Heart Foundation of Australia and Kinect Australia for the Lifescripts Consortium (2005) Lifescripts in your division: supporting lifestyle risk factor management in general practice. In *A guide for Divisions of General Practice*. Commonwealth of Australia, Canberra.
- Newell, S., Girgis, A., Sanson-Fisher, R. and Savolainen, N. (1999) The accuracy of self-reported health behaviors and risk factors relating to cancer and cardiovascular disease in general population: a critical review. *American Journal of Preventive Medicine*, **17**, 211–229.
- Rogers, C. (1951) *Client-centered Therapy: Its Current Practice, Implications and Theory*. Houghton Mifflin, Boston.
- Roter, D. (1991) *Roter Interaction Analysis Coding System (RIAS), Coding Manual*. John Hopkins University, Baltimore.
- Sciamanna, C. N., DePue, J. D., Goldstein, M. G., Park, E. R., Gans, K. M., Monroe, A. D. *et al.* (2002) Nutrition counseling in the Promoting Cancer Prevention in Primary Care Study. *Preventive Medicine*, **35**, 437–446.
- Serdula, M. K., Khan, L. K. and Dietz, W. H. (2003) Weight loss counseling revisited. *Journal of the American Medical Association*, **289**, 1747–1750.
- Shi, L., Starfield, B. and Xu, J. (2001) Validating the adult primary care assessment tool. *The Journal of Family Practice*, **50**, 161W–175W.
- Stange, K. C., Zyzanski, S. J., Smith, T. F., Kelly, R., Langa, D. M., Flocke, S. A., *et al.* (1998) How valid are medical records and patient questionnaires for physician profiling and health services research? *Medical Care*, **36**, 851–857.
- Stange, K. C., Flocke, S. A. G. M. A., Kelly, R. B. and Zyzanski, S. J. (2000) Direct observation of rates of preventive service delivery in community family practice. *Preventive Medicine*, **31**, (2 Pt 1), 167–176.
- The Quality Indicator Study Group (1995) An approach to the evaluation of quality indicators of the outcome of care in hospitalized patients, with a focus on nosocomial infection indicators. *Infection Control and Hospital Epidemiology*, **16**, 308–316.
- The Royal Australian College of General Practitioners (2004) A population health guide to behavioral risk factors in general practice. In *Smoking, nutrition, alcohol and physical activity (SNAP)* (<http://www.racgp.org.au/document.asp?id=14803>).
- Truax, C. B. (1966) Therapists' empathy, warmth, genuineness, and patient personality change in group psychotherapy. *Journal of Clinical Psychology*, **22**, 225–229.
- Wagner, E. H. (1998) Chronic disease management: what will it take to improve care for chronic illness. *Effective Clinical Practice*, **1**, 1–4.